

Note: Handouts DO NOT replace the book. In most cases, they only provide a guideline on topics and an intuitive feel. The math details will be covered in class, so it is important to attend class and also you MUST read the book.

1 Motivation and Applications

- As stated by Laplace, “Probability is common sense reduced to calculation”.
- You need to first learn the theory required to correctly do these calculations. The examples that I solve and those in the book and the homeworks will provide a wonderful practical motivation as to why you need to learn the theory.
- If you patiently grasp the basics, especially the first 4 chapters of BT, it will be the most useful thing you’ve ever learnt - whether you pursue a career in EE or CE or Economics or Finance or Management and also while you try to invest in stocks or gamble in Las Vegas!
- Applications: communications (telephones, cell phones, TV, ...), signal processing (image and video denoising, face recognition, tracking moving objects from video,...), systems and control (operating an airplane, fault detection in a system,...), predicting reliability of a system (e.g. electrical system), resource allocation, internet protocols, non-engineering applications (e.g. medicine: predicting how prevalent a disease is or well a drug works, weather forecasting, economics).

2 Introduction: Topics Covered (BT, 1.1 - 1.6)

- What is Probability
- Set Theory Basics
- Probability Models
- Conditional Probability
- Total Probability and Bayes Rule
- Independence
- Counting

3 What is Probability?

- Measured relative frequency of occurrence of an event.
Example: toss a coin 100 times, measure frequency of heads or compute probability of raining on a particular day and month (using past years’ data)
- Or subjective belief about how “likely” an event is (when do not have data to estimate frequency).
Example: any one-time event in history or “how likely is it that a new experimental drug will work?”
- Two types of problems need to be solved

1. Specify the probability “model” or learn it (statistics).
2. Use the “model” to compute probability of different events.

- **We will assume the model is given and will focus on 2. in this course.**

4 Set Theory Basics

- Set: any collection of objects (elements of a set).
- Discrete sets
 - Finite number of elements, e.g. numbers of a die
 - Or infinite but countable number of elements, e.g. set of integers
- Continuous sets
 - Cannot count the number of elements, e.g. all real numbers between 0 and 1.
- “Universe” (denoted Ω): consists of all possible elements that could be of interest. In case of random experiments, it is the set of all possible outcomes. Example: for coin tosses, $\Omega = \{H, T\}$.
- Empty set (denoted ϕ): a set with no elements
- Subset: $A \subseteq B$: if every element of A also belongs to B.
- Strict subset: $A \subset B$: if every element of A also belongs to B and B has more elements than A.
- Belongs: \in , Does not belong: \notin
- Complement: A' or A^c , Union: $A \cup B$, Intersection: $A \cap B$
 - $A' \triangleq \{x \in \Omega | x \notin A\}$
 - $A \cup B \triangleq \{x | x \in A, \text{ or } x \in B\}$, $x \in \Omega$ is assumed.
 - $A \cap B \triangleq \{x | x \in A, \text{ and } x \in B\}$
 - Visualize using Venn diagrams (see book)
- **Disjoint sets: A and B are disjoint if $A \cap B = \phi$ (empty), i.e. they have no common elements.**
- DeMorgan’s Laws

$$(A \cup B)' = A' \cap B' \tag{1}$$

$$(A \cap B)' = A' \cup B' \tag{2}$$

- Proofs: Need to show that every element of LHS (left hand side) is also an element of RHS (right hand side) AND vice versa. Details in class.
- Read section on Algebra of Sets, pg 5

5 Probabilistic models

- There is an underlying process called **experiment** that produces exactly ONE **outcome**.
- A probabilistic model: consists of a sample space and a probability law
 - Sample space (denoted Ω): set of all possible outcomes of an experiment
 - Event: any subset of the sample space
 - Probability Law: assigns a probability to every set A of possible outcomes (event)
 - Choice of sample space (or universe): every element should be distinct and mutually exclusive (disjoint); and the space should be “collectively exhaustive” (every possible outcome of an experiment should be included).
- **Probability Axioms:**
 1. **Nonnegativity.** $P(A) \geq 0$ for every event A .
 2. **Additivity.** If A and B are two **disjoint** events, then $P(A \cup B) = P(A) + P(B)$ (also extends to any countable number of disjoint events).
 3. **Normalization.** Probability of the entire sample space, $P(\Omega) = 1$.
- Probability of the empty set, $P(\phi) = 0$ (follows from Axioms 2 & 3).
- Sequential models, e.g. three coin tosses or two sequential rolls of a die.
Tree-based description: see Fig. 1.3
- Discrete probability law: sample space consists of a finite number of possible outcomes, law specified by probability of single element events.
 - Example: for a fair coin toss, $\Omega = \{H, T\}$, $P(H) = P(T) = 1/2$
 - Discrete uniform law for any event A : $P(A) = \frac{\text{number of elements in } A}{n}$
- Continuous probability law: e.g. $\Omega = [0, 1]$: probability of any single element event is zero, need to talk of probability of a subinterval, $[a, b]$ of $[0, 1]$.
Example 1.4, 1.5
- Properties of probability laws
 1. If $A \subseteq B$, then $P(A) \leq P(B)$
 2. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 3. $P(A \cup B) \leq P(A) + P(B)$
 4. $P(A \cup B \cup C) = P(A) + P(A' \cap B) + P(A' \cap B' \cap C)$
 5. Note: book uses A^c for A' (complement of set A).
 6. Proofs: in book. Some will be done in class.
 7. Visualize: see Venn diagrams in book.

6 Conditional Probability

- Given that we know that an event B has occurred, what is the probability that event A occurred? Denoted by $P(A|B)$. Example: Roll of a 6-sided die. Given that the outcome is even, what is the probability of a 6? Answer: $1/3$
- When number of outcomes is finite and all are equally likely,

$$P(A|B) = \frac{\text{number of elements of } A \cap B}{\text{number of elements of } B} \quad (3)$$

- In general,

$$P(A|B) \triangleq \frac{P(A \cap B)}{P(B)} \quad (4)$$

- $P(A|B)$ is a probability law (satisfies axioms) on the universe B . Exercise: show this.
- Examples/applications
 - Example 1.7, 1.8, 1.11
 - Construct sequential models: $P(A \cap B) = P(B)P(A|B)$. Example: Radar detection (Example 1.9). What is the probability of the aircraft not present and radar registers it (false alarm)?
 - See Fig. 1.9: Tree based sequential description

7 Total Probability and Bayes Rule

- Total Probability Theorem: Let A_1, \dots, A_n be disjoint events which form a partition of the sample space ($\cup_{i=1}^n A_i = \Omega$). Then for any event B ,

$$\begin{aligned} P(B) &= P(A_1 \cap B) + \dots + P(A_n \cap B) \\ &= P(A_1)P(B|A_1) + \dots + P(A_n)P(B|A_n) \end{aligned} \quad (5)$$

Visualization and proof: see Fig. 1.13

- Example 1.13, 1.15
- Bayes rule: Let A_1, \dots, A_n be disjoint events which form a partition of the sample space. Then for any event B , s.t. $P(B) > 0$, we have

$$P(A_i|B) = \frac{P(A_i)P(B|A_i)}{P(B)} = \frac{P(A_i)P(B|A_i)}{P(A_1)P(B|A_1) + \dots + P(A_n)P(B|A_n)} \quad (6)$$

- Inference using Bayes rule
 - There are multiple “causes” A_1, A_2, \dots, A_n that result in a certain “effect” B . Given that we observe the effect B , what is the probability that the cause was A_i ? Answer: use Bayes rule. See Fig. 1.14
 - Radar detection: what is the probability of the aircraft being present given that the radar registers it? Example 1.16
 - False positive puzzle, Example 1.18: very interesting!

8 Independence

- $P(A|B) = P(A)$ and so $P(A \cap B) = P(B)P(A)$: the fact that B has occurred gives no information about the probability of occurrence of A.
Example: A = head in first coin toss, B = head in second coin toss.
- “Independence”: DIFFERENT from “mutually exclusive” (disjoint)
 - Events A and B are disjoint if $P(A \cap B) = 0$: cannot be independent if $P(A) > 0$ and $P(B) > 0$.
Example: A = head in a coin toss, B = tail in a coin toss
 - Independence: a concept for events in a sequence. Independent events with $P(A) > 0, P(B) > 0$ cannot be disjoint
- Conditional independence **
- Independence of a collection of events
 - $P(\cap_{i \in S} A_i) = \prod_{i \in S} P(A_i)$ for every subset S of $\{1, 2, \dots, n\}$
- Reliability analysis of complex systems: independence assumption often simplifies calculations
 - Analyze Fig. 1.15: what is $P(\text{system fails})$ of the system $A \rightarrow B$?
 - * Let p_i = probability of success of component i .
 - * m components in series: $P(\text{system fails}) = 1 - p_1 p_2 \dots p_m$ (succeeds if all components succeed).
 - * m components in parallel: $P(\text{system fails}) = (1 - p_1) \dots (1 - p_m)$ (fails if all the components fail).
- Independent Bernoulli trials and Binomial probabilities
 - A Bernoulli trial: a coin toss (or any experiment with two possible outcomes, e.g. it rains or does not rain, bit values)
 - Independent Bernoulli trials: sequence of independent coin tosses
 - Binomial: Given n independent coin tosses, what is the probability of k heads (denoted $p(k)$)?
 - * probability of any one sequence with k heads is $p^k(1 - p)^{n-k}$
 - * number of such sequences (from counting arguments): $\binom{n}{k}$
 - * $p(k) = \binom{n}{k} p^k(1 - p)^{n-k}$, where $\binom{n}{k} \triangleq \frac{n!}{(n-k)!k!}$
 - Application: what is the probability that more than c customers need an internet connection at a given time? We know that at a given time, the probability that any one customer needs connection is p .
Answer:
$$\sum_{k=c+1}^n p(k)$$

9 Counting

- Needed in many situations. Two examples are:
 1. Sample space has a finite number of equally likely outcomes (discrete uniform), compute probability of any event A.
 2. Or compute the probability of an event A which consists of a finite number of equally likely outcomes each with probability p , e.g. probability of k heads in n coin tosses.
- Counting principle (See Fig. 1.17): Consider a process consisting of r stages. If at stage 1, there are n_1 possibilities, at stage 2, n_2 possibilities and so on, then the total number of possibilities = $n_1 n_2 \dots n_r$.
 - Example 1.26 (number of possible telephone numbers)
 - Counting principle applies even when second stage depends on the first stage and so on, Ex. 1.28 (no. of words with 4 distinct letters)
- Applications: k -permutations.
 - n distinct objects, how many different ways can we pick k objects and arrange them in a sequence?
 - * Use counting principle: choose first object in n possible ways, second one in $n - 1$ ways and so on. Total no. of ways: $n(n - 1) \dots (n - k + 1) = \frac{n!}{(n-k)!}$
 - * If $k = n$, then total no. of ways = $n!$
 - * Example 1.28, 1.29
- Applications: k -combinations.
 - Choice of k elements out of an n -element set without regard to order.
 - Most common example: There are n people, how many different ways can we form a committee of k people? Here order of choosing the k members is not important. Denote answer by $\binom{n}{k}$
 - Note that selecting a k -permutation is the same as first selecting a k -combination and then ordering the elements (in $k!$) different ways, i.e. $\frac{n!}{(n-k)!} = \binom{n}{k} k!$
 - Thus $\binom{n}{k} = \frac{n!}{k!(n-k)!}$.
 - How will you relate this to the binomial coefficient (number of ways to get k heads out of n tosses)?
Toss number j = person j , a head in a toss = the person (toss number) is in committee
- Applications: k -partitions. **
 - A combination is a partition of a set into two parts
 - Partition: given an n -element set, consider its partition into r subsets of size n_1, n_2, \dots, n_r where $n_1 + n_2 + \dots + n_r = n$.
 - * Use counting principle and k -combinations result.
 - * Form the first subset. Choose n_1 elements out of n : $\binom{n}{n_1}$ ways.
 - * Form second subset. Choose n_2 elements out of $n - n_1$ available elements: $\binom{n - n_1}{n_2}$ and so on.
 - * Total number of ways to form the partition: $\binom{n}{n_1} \binom{n - n_1}{n_2} \dots \binom{n - n_1 - n_2 - \dots - n_{r-1}}{n_r} = \frac{n!}{n_1! n_2! \dots n_r!}$